



Protein Analyzer

AR-100



AR-100



Reagent Kits

Item	Packing	Speciment type	Test Range	Reference Range	Clinical Significance
CRP	25 Test 50 Test	5µl Serum	0-180 mg/L	< 6 mg/L	Identify bacterial infection or Virus infection. Dynamically monitor the course of disease. Observe the effectiveness of antibiotics.
hs-CRP	25 Test 50 Test	20µl serum	0,20-16,0 mg/l	0,1-10 mg/l	Predict morbidity and mortality of cardiovascular disease. Identify newborn bacterial infection and guide clinical use of antibiotics.
mAlb	25 Test 50 Test	100µl urine	10-160mg/l	<20mg/l	Assess the early lesions of kidney. An ideal endogenous marker of the glomerular filtration rate. Monitor the early lesions of diabetic Mellitus. Discover the possibilities of cardiovascular accident and unhealthy kidney function of hypertensive patient.
Cys-c	25 Test 50 Test	10µl Serum	0,2-6,0 mg/l	<1,45mg/l	Monitor the early lesions of kidney caused by urinary tract infection etc.
RA	25 Test 50 Test	5µl Serum	0-140 IU/ML	< 20 IU/ML	Rheumatoid factor is a group of antibodies directed to determinants in the Fc portion of the immunoglobulin G-molecule.
ASO	25 Test 50 Test	5µl Serum	0-500 IU/ml	< 200 IU/ml	SLO is a toxic immunogenic coenzyme produced by b-hemolytic streptococci of group A,C and G. Measuring the ASO antibodies are useful for the diagnosis of rheumatoid fever, acute glomerulonephritis and streptococcal infections.
HBA1C	25 Test 50 Test	5µl Heamolysate	0-18,26%	< 6 %	HBA1C is a glycated Product of Hemoglobin AO (HBA1C), The Predominant form of hemoglobin in adult.
Vitamin D	25 Test	5µl Serum	5-130 ng/ml	30-100 ng/ml	Vitamin D Helps your body absorb Calcium and maintain Strong bones throughout your entire life.
Anti CCP	25 Test	5µl Serum	25-3200 U/ml	<25 U/ml	Anti-Cyclic Citrullinated peptide (CCP) antibody has been reported to greatly improve the diagnostic efficiency of RA.
D-dimer	25 Test	20µl Serum	0-10.8 ng/ml	<25 U/ml	Increase of D-dimer in blood testifies the blood clot is formed and fibrinolytic activity has functioned. It is known that high value of D-dimer is indicated in disease such as malignant tumor, vascular disease.
Ferritin	25 Test	10µl Serum	20-1000 ng/ml	20-250 ng/ml	The determination of Serum Ferritin concentration is very useful for the diagnosis, treatment and prognosis of iron metabolism abnormalities such as anaemia and iron excess, liver disease etc.

SPECIFICATIONS

Specimen type	Peripheral whole blood, vein whole blood, serum, plasma, urine specimens
Specimen No	Automatic numbering 1 -999 for the same test, optional barcode scanner
Whole blood HCT	00.00-99.99% adjustable
Specimen dilution	1-50 times adjustable;
Test speed	20-120 sec/test;
Screen	5.5 inch LCD touch screen
Interface	3 RS232C interface (Data transmission, external printer, external barcode scanner)
Result storage	10000
Operating voltage	AC 100V-240V/DC 12V



Service Helpline
1800-889-4232

ACER BIOMEDICALS LLP

Office: Industrial unit -80, Block-G, Sector-63, Noida, U.P.(India)

+91 120 4166921, E-mail: info@acerbiomed.com

Web: www.acerbiomedicals.com, www.acerbiomed.com